arXiv:1003.1034v1 [math.GT] 4 Mar 2010

RECURRENCE RELATION FOR HOMFLY POLYNOMIAL AND RATIONAL SPECIALIZATIONS

REHANA ASHRAF[1], BARBU BERCEANU[1,2]

Abstract. Turning the skein relation for HOMFLY into a Fibonacci recurrence, we prove that there are only three rational specializations of HOMFLY polynomial: Alexander-Conway, Jones, and a new one. Using the recurrence relation, we find general and relative expansion formulae and rational generating functions for Alexander-Conway polynomial and the new polynomial, which reduce the computations to closure of simple braids, a subset of square free braids; HOMFLY polynomials of these simple braids are also computed. Algebraic independence of these three polynomials is proved.

1. **Introduction**

HOMFLY polynomial (see [6], [8]) is a function P : {oriented link diagrams} $\longrightarrow$ $\mathbb{C}[l^{\pm 1}, m^{\pm 1}]$ satisfying:

a) **Normalization:** $P(\bigcirc) = 1$;

b) **Skein Relation:** whenever three oriented link diagrams L_+, L_- and L_0 are the same, except in the neighborhood of one crossing, where



then

$$lP(L_+) + l^{-1}P(L_-) + mP(L_0) = 0.$$

HOMFLY polynomial is in fact a link invariant which generalizes both Alexander-Conway and Jones polynomials. The specializations:

---

keywords and phrases: HOMFLY polynomial, Alexander-Conway polynomial, simple braids, Fibonacci recurrence.

This research is partially supported by Higher Education Commission, Pakistan.

2010 AMS classification: Primary 57M27, 57M25; Secondary 20F36.

1) $l = \iota t^{-1}, m = \iota(t^{-\frac{1}{2}} - t^{\frac{1}{2}})$ gives the Jones polynomial $V(L)$ and skein relation

$$tV(L_-) - (t^{-\frac{1}{2}} - t^{\frac{1}{2}})V(L_0) - t^{-1}V(L_+) = 0, \text{ respectively}$$

2) $l = \iota, m = \iota(t^{\frac{1}{2}} - t^{-\frac{1}{2}})$ gives the Alexander-Conway polynomial $\nabla(L)$ and skein relation

$$\nabla(L_-) - (t^{\frac{1}{2}} - t^{-\frac{1}{2}})\nabla(L_0) - \nabla(L_+) = 0.$$

In this paper we transform the skein relation into a Fibonacci type recurrence for HOMFLY polynomial of closed braids. The generators $x_i\,(i = 1, \ldots, n-1)$ of the braid group $\mathcal{B}_n$ are classical Artin generators



The braid group $\mathcal{B}_n$ is fixed, and also the sequence of generators $(x_{i_1}, x_{i_2}, \ldots, x_{i_k})$. Let us denote by $P_n(a_1, \ldots, a_k)$ HOMFLY polynomial of the closure of the n-braid $\beta = x_{i_1}^{a_1} x_{i_2}^{a_2} \ldots x_{i_k}^{a_k}$. The first result of the paper is

Theorem 1.1. *For any $a_1, \ldots, a_k \in \mathbb{Z}$ and any $j = 1, \ldots, k$ the recurrence holds*

$$P_n(a_1, \ldots, a_j+2, \ldots, a_k) + mlP_n(a_1, \ldots, a_j+1, \ldots, a_k) + l^2 P_n(a_1, \ldots, a_j, \ldots, a_k) = 0.$$

Using the terminology of [9], the sequence $\{P_n(a_1, \ldots, a_k)\}_{a_* \in \mathbb{Z}^k}$ is *a multiple Fibonacci sequence* with parameters $(-ml, -l^2)$. Make the change of variable $t \to s^{-2}$, Alexander-Conway and Jones specializations give two recurrences; in the next formulae $\nabla_n(a_1, \ldots, a_k)$, $V_n(a_1, \ldots, a_k)$ (and also $D_n(a_1, \ldots, a_k)$) stand for corresponding polynomials of the closure of n-braid $\beta = x_{i_1}^{a_1} x_{i_2}^{a_2} \ldots x_{i_k}^{a_k}$.

Corollary 1.2. *Alexander-Conway polynomial satisfies the multiple Fibonacci recurrence:*

$$\nabla_n(a_1, \ldots, a_j+2, \ldots, a_k) = (s^{-1}-s)\nabla_n(a_1, \ldots, a_j+1, \ldots, a_k) + \nabla_n(a_1, \ldots, a_j, \ldots, a_k).$$

Corollary 1.3. (see [3]) *Jones polynomial satisfies the multiple Fibonacci recurrence:*

$$V_n(a_1, \ldots, a_j+2, \ldots, a_k) = (s^3-s)V_n(a_1, \ldots, a_j+1, \ldots, a_k) + s^4 V_n(a_1, \ldots, a_j, \ldots, a_k).$$

A "relative" example of multiple Fibonacci sequence is given by HOMFLY polynomials of the link diagrams $\mathcal{L}(R_{k_1}, \ldots, R_{k_p})$, where there are p subdiagrams and the diagram with index j contains the braid $x_1^{k_j}$, ($k_1, \ldots, k_p$ are arbitrary integers)



and the rest of the diagram is fixed. HOMFLY polynomials of this family, denoted by $P_{\mathcal{L}}(k_1,\dots,k_p)$, satisfy the same multiple Finonacci recurrence:

Proposition 1.4. *For any $k_1,\dots,k_p$ and $j\in\{1,2,\dots,p\}$*

$$P_{\mathcal{L}}(k_1,\dots,k_j+2,\dots,k_p)=-mlP_{\mathcal{L}}(k_1,\dots,k_j+1,\dots,k_p)-l^2P_{\mathcal{L}}(k_1,\dots,k_j,\dots,k_p).$$

The roots of characteristic equations corresponding to these recurrences are $r_1=s^{-1}$, $r_2=-s$ for Alexander-Conway and $r_1'=s^3$, $r_2'=-s$ for Jones [3]. It turns out that, up to constants, Alexander-Conway and Jones specializations are unique with the property that the roots of the characteristic equation are s and $s^m, m\in\mathbb{Z}$, $m\neq 1$; the case of double roots $r_1''=r_2''=s$ gives a new polynomial (see Theorem 2.4).

Section 2 of the paper contains the proof of Theorem 1.1 and precise definitions of rational and normalized specializations (see Definition 2.2 and Definition 2.3).

In the next section two direct consequences of the recurrence relation for Alexander-Conway polynomial are given. The module of multiple Fibonacci sequences is a tensor power of a free module (see [9]), therefore one can describe such sequences using a canonical basis: let us introduce the Laurent polynomial

$$C^a(s)=(-1)^a s^{a-1}+s^{1-a}.$$

Theorem 1.5. ***(Expansion formula)*** *The Alexander-Conway polynomial of the closure of the n-braid $\beta=x_{i_1}^{a_1}x_{i_2}^{a_2}\dots x_{i_k}^{a_k}$ is given by*

$$\nabla_n(x_{i_1}^{a_1}\dots x_{i_k}^{a_k})(s)=\Big(\frac{s}{s^2+1}\Big)^k\sum_{0\le j_1,\dots,j_k\le 1}C^{a_1+j_1}(s)\dots C^{a_k+j_k}(s)\nabla_n(x_{i_1}^{j_1}\dots x_{i_k}^{j_k})(s).$$

Practitioners of Burau representation of braid groups could compare the complexity of computation of Burau matrix and its determinant with the expansion formula. People working with Pretzel links and 2-bridge knots could find the next result familiar:

Corollary 1.6. ***(Relative expansion formula)*** *The Alexander-Conway polynomial of the link* $\mathcal{L}(R_{k_1},\dots,R_{k_p})$ *is*

$$\nabla_{\mathcal{L}}(k_1,\dots,k_p)=\Big(\frac{s}{s^2+1}\Big)^p\sum_{0\le j_1,\dots,j_p\le 1}C^{k_1+j_1}(s)\dots C^{k_p+j_p}(s)\nabla_{\mathcal{L}}(j_1,\dots,j_p)(s).$$

For the next result n and the sequence $(x_{i_1},\dots,x_{i_k})$ of generators of $\mathcal{B}_n$ are fixed. In order to compute the *generating function* of Alexander-Conway polynomials of type $I_*=(i_1,\dots,i_k)$, the formal series in $t_1,\dots,t_k$:

$$\mathcal{AC}_n(t_1,\dots,t_k)=\sum_{(a_1,a_2,\dots,a_k)\in\mathbb{Z}^k}\nabla_n(x_{i_1}^{a_1}\dots x_{i_k}^{a_k})(s)t_1^{a_1}\dots t_k^{a_k},$$

we introduce new polynomials in the formal variable τ :

$$q(\tau)=(1+s\tau)(1-s^{-1}\tau),\ Q_0(\tau)=1-(s^{-1}-s)\tau \text{ and } Q_1(\tau)=\tau.$$

Theorem 1.7. ***(Generating function)*** *The generating function* $\mathcal{AC}_n(t_1,\dots,t_k)$ *for Alexander-Conway polynomials is a rational function in* $t_1.\dots,t_k$*:*

$$\mathcal{AC}_n(t_1,\dots,t_k)=q(t_1)^{-1}\dots q(t_k)^{-1}\sum_{0\le j_1,\dots,j_k\le 1}Q_{j_1}(t_1)\dots Q_{j_k}(t_k)\nabla_n(x_{i_1}^{j_1}\dots x_{i_k}^{j_k}).$$

This section contains also some computations of Alexander-Conway polynomial.

In section 4 the degenerate case $(r_1=r_2=s)$ is analyzed. We find a new polynomial D satisfying:

a) **Normalization:** $D(\bigcirc)=1$;

b) **Skein Relation**: $sD(L_+)+s^{-1}D(L_-)-2D(L_0)=0$,

and corresponding expansion formulae

Theorem 1.8. *(**Expansion formula**) The* D *polynomial of the closure of the* n*-braid* $\beta=x_{i_1}^{a_1}x_{i_2}^{a_2}\dots x_{i_k}^{a_k}$ *is*

$$D_n(x_{i_1}^{a_1}x_{i_2}^{a_2}\dots x_{i_k}^{a_k})(s)=\sum_{0\le j_1,\dots,j_k\le 1}R_{j_1}^{[a_1]}(s)\dots R_{j_k}^{[a_k]}(s)D_n(x_{i_1}^{j_1}\dots x_{i_k}^{j_k})(s).$$

(in the previous and next formulae: $R_0^{[a]}(s)=(1-a)s^a$, $R_1^{[a]}(s)=as^{a-1}$),

Corollary 1.9. ***(Relative expansion formula)*** *The* D *polynomial of the link* $\mathcal{L}(R_{k_1},\dots,R_{k_p})$ *is*

$$D_{\mathcal{L}}(k_1,\dots,k_p)=\sum_{0\le j_1,\dots,j_p\le 1}R_{j_1}^{[k_1]}(s)\dots R_{j_p}^{[k_p]}D_{\mathcal{L}}(j_1,\dots,j_p)(s).$$

and a rational *generating function*

Theorem 1.10. ***(Generating function)*** *The generating function* $\mathcal{D}_n(t_1,\ldots,t_k)$ *for the degenerate polynomials* $D_n(x_{i_1}^{a_1}x_{i_2}^{a_2}\ldots x_{i_k}^{a_k})$ *is a rational function in* $t_1,\ldots,t_k$

$$\mathcal{D}_n(t_1,\ldots,t_k)=e(t_1)^{-1}\ldots e(t_k)^{-1}\sum_{0\le j_1,\ldots,j_k\le 1}E_{j_1}(t_1)\ldots E_{j_k}(t_k)D_n(x_{i_1}^{j_1}\ldots x_{i_k}^{j_k})$$

where $e(\tau)=(1-s\tau)^2$, $E_0(\tau)=1-s\tau$, $E_1(\tau)=\tau$.

Some computations of D polynomials are given at the end of section 4.

The expansion formula of Theorems 1.5 and 1.8 reduce the computations of link polynomials to polynomials of positive braids with total exponent $J=j_1+j_2+\ldots+j_k\le k$. If these braids (or some positive conjugates of them) contain exponents greater than 1, apply again and again expansion formula; finally we obtain *simple braids.* In section 5 we give different characterizations of this set of braids, the proofs and more properties are contained in [2]. The first phrase in the next theorem could be taken as a definition :

Theorem 1.11. *a) A positive braid in* $\mathcal{B}_n$ *is simple if and only if it is conjugate to*

$$\beta_A=(x_1x_2\ldots x_{s_1-1})(x_{s_1+1}\ldots x_{s_2-1})\ldots(x_{s_{r-1}+1}\ldots x_{s_r-1})$$

for an decreasing sequence $A=(a_1,a_2,\ldots,a_r)$ *with* $a_1\ge a_2\ge\ldots\ge a_r\ge 2$ *and* $s_i=a_1+a_2+\ldots+a_i$.
b) HOMFLY polynomial of a closed simple braid B_A *is given by*

$$P_n(\beta_A)(l,m)=\Big(-\frac{l+l^{-1}}{m}\Big)^{n-s_r+r-1}.$$

The values of Alexander-Conway, Jones and D polynomials for simple braids are computed in section 5.

In the last section we show that Alexander-Conway, Jones and D polynomials of 2-braids are linearly independent over $\mathbb{C}[s,s^{-1}]$ and also

Theorem 1.12. *The link polynomials* ∇, V *and* D *are algebraically independent over* $\mathbb{C}[s,s^{-1}]$.

The last example is a 2-link with the same Alexander-Conway and Jones polynomials as the unlink, but with a nontrivial D invariant.

2. Nice Specializations

Proof of Theorem 1.1: Consider the braid (the exponent e is positive)

$$\beta(e+2)=x_{i_1}^{a_1}x_{i_2}^{a_2}\ldots x_{i_{j-1}}^{a_{j-1}}x_{i_j}^{e+2}x_{i_{j+1}}^{a_{j+1}}\ldots x_{i_k}^{a_k}.$$

Applying the skein relation for HOMFLY polynomial of this braid, the geometrical change appears as a change at the last crossing of the j-th position. Both parts $x_{i_1}^{a_1}x_{i_2}^{a_2}\ldots x_{i_{j-1}}^{a_{j-1}}$ and $x_{i_{j+1}}^{a_{j+1}}\ldots x_{i_{k-1}}^{a_{k-1}}x_{i_k}^{a_k}$ remain unchanged. The local picture is:



The skein relation $lP(L_+) + l^{-1}P(L_-) + mP(L_0) = 0$ with correspondence $L_- \to \widehat{\beta}(e+2)$, $L_0 \to \widehat{\beta}(e+1)$, $L_+ \to \widehat{\beta}(e)$, gives

$$P_n(e+2) + mlP_n(e+1) + l^2P_n(e) = 0.$$

The case $e < 0$ can be reduced to the above case by adding a new factor on the $j+1$ position, $x_{i_1}^{a_1}\ldots x_{i_j}^{e}\ldots x_{i_k}^{a_k} = x_{i_1}^{a_1}\ldots x_{i_j}^{d+e}x_{i_j}^{-d}\ldots x_{i_k}^{a_k}$, with d big enough.

□

Example 2.1. A more complicated recurrence is used for the sequence $P(k) = P_3(\gamma_k)$ where γ_k is the 3-braid $x_1x_2x_1\ldots(k-\text{factors})$. This sequence contains the powers of Garside braid Δ_3: $\gamma_{6k} = \Delta_3^{2k}$, $\gamma_{6k+3} = \Delta_3^{2k+1}$.

1) $P(2k+1) = -mlP(2k) - l^2P(2k-1)$
2) $P(6k+4) = -mlP(6k+3) - l^2P(6k+2)$
3) $P(6k+2) = -mlP(6k+1) + ml^3P(6k-1) + l^4P(6k-2)$
4) $P(6k) = -mlP(6k-1) + ml^3P(6k-3) - ml^5P(6k-5) - l^6P(6k-6)$.

The proof is the same as in [3]. The characteristic polynomial of this recurrence has 0 a triple root. In the case of Jones specialization the other three roots are s^6, s^{12} and s^{18}. For Alexander-Conway polynomial, see section 3 (and for D polynomial see section 4).

The roots of the characteristic equation of the basic recurrence of Theorem 1.1 are not rational functions in l, m. From the point of view of recurrence relations a "*nice specialization*" $l \to F(s)$, $m \to G(s)$ will give "*nice roots*" of the characteristic equation.

Definition 2.2. Let us call a *rational specialization* of HOMFLY $P(l, m)$ a specialization $l = F(s)$, $m = G(s)$ (F, G, are rational functions) such that:
a) $P(\bigcirc\bigcirc)(F(s), G(s))$ is a Laurent polynomial in s;
b) the roots of the characteristic equation are $r_1 = \lambda s^n$, $r_2 = \mu s^k$ where n, k are coprime integers and λ, μ are non zero complex numbers.

Definition 2.3. A rational specialization is *normalized* if n equals 1 and *nondegenerate* if $n \neq k$.

The next result says that there are few "nice specializations" and also that Laurent condition *a)* on the unlink gives Laurent polynomials for any link.

Theorem 2.4. *There are (up to constants) only three normalized rational specialization of the HOMFLY polynomial, two nondegenerate and one degenerate:*
1) Alexander-Conway with roots $r_1 = -s$, $r_2 = s^{-1}$*;*
2) Jones with roots $r_1 = -s$, $r_2 = s^3$;
3) roots $r_1 = r_2 = s$.

Proof. The characteristic equation of the HOMFLY recurrence is

$$r^2 + mlr + l^2 = 0.$$

Let $\lambda^2 s^n$, $\mu^2 s^k$ be the roots of the equation, where $\lambda^2, \mu^2 \in \mathbb{C}^*$, $n, k \in \mathbb{Z}$, $(n; k) = 1$. We should have $n + k = 2q$ for some integer q, therefore

$$(*) \qquad l \;=\; \lambda\mu s^q,\; m = -\frac{\lambda}{\mu}s^{n-q} - \frac{\mu}{\lambda}s^{q-n}.$$

Skein relation for eight figure ∞ gives:

$$\begin{aligned} P(\bigcirc\bigcirc) &= -\frac{l + l^{-1}}{m} = \frac{\lambda\mu s^q + (\lambda\mu s^q)^{-1}}{\lambda\mu^{-1}s^{n-q} + \mu\lambda^{-1}s^{q-n}} \\ &= \frac{\lambda^2\mu^2 s^{2q} + 1}{\lambda\mu s^q}\frac{\lambda\mu s^{n-q}}{\lambda^2 s^{2(n-q)} + \mu^2} = \frac{1}{s^k}\cdot\frac{\lambda^2\mu^2 s^{2q} + 1}{\lambda^2 s^{2(n-q)} + \mu^2} \end{aligned}$$

and the first condition for a rational parametrization implies $n - q \mid q$ or $n - q = 0$. The case when $n - q = 0$ gives the degenerate case $n = k = 1$, $r_1 = r_2 = s$. If $n - q \mid q$, using $(n; k) = 1$, we have $(n; q) = 1$, hence $n - q = \pm 1$. Normalization implies $n = 1$, hence $q = 0$, $k = -1$ or $q = 2$, $k = 3$. One can find Alexander-Conway and Jones specializations (up to constants) taking these values for n and q in $(*)$. $\square$

Fibonacci recurrence for Jones polynomial is treated in [3]. In the next two sections we analyze the recurrence for Alexander-Conway and for the degenerate case.

3. Alexander-Conway polynomial

Using Alexander-Conway specialization, $l = \iota, m = \iota(s^{-1} - s)$, the recurrence relation for Alexander-Conway polynomials is (Corollary 1.2)

$$\nabla_n(a_1, \ldots, a_j{+}2, \ldots, a_k) = (s^{-1}{-}s)\nabla_n(a_1, \ldots, a_j{+}1, \ldots, a_k){+}\nabla_n(a_1, \ldots, a_j, \ldots, a_k).$$

The characteristic equation $r^2 = (s^{-1} - s)r + 1$ has the roots $r_1 = -s$, $r_2 = s^{-1}$. Applying computations with multiple Fibonacci sequence (see [9], Theorem 3.8, for complete details) one can find the expansion formula and compute the generating function.

Proof of Theorem 1.5: Laurent polynomials of ([9], Theorem 3.8 a) are given by:

$$\begin{aligned}
\Delta &= r_2 - r_1 = \frac{s^2+1}{s} \\
S_0^{[a]} &= r_1^a r_2 - r_1 r_2^a = (-1)^a s^{a-1} + s^{1-a} = C^a(s) \\
S_1^{[a]} &= r_2^a - r_1^a = s^{-a} + (-1)^{a+1} s^a = C^{a+1}(s),
\end{aligned}$$

therefore the general term is given by

$$\nabla_n(x_{i_1}^{a_1} \ldots x_{i_k}^{a_k})(s) = \left(\frac{s}{s^2+1}\right)^k \sum_{0 \le j_1, \ldots, j_k \le 1} C^{a_1+j_1}(s) \ldots C^{a_k+j_k}(s) \nabla_n(x_{i_1}^{j_1} \ldots x_{i_k}^{j_k})(s).$$

□

Proof of Theorem 1.7 From the second part of ([9], Theorem 3.8), we use the polynomials of formal variables of $t_1, t_2, \ldots, t_k$ (with coefficients in the ring of Laurent polynomials in variable s):

$$\begin{aligned}
q(\tau) &= (1 - r_1\tau)(1 - r_2\tau) = 1 + (s - s^{-1})\tau - \tau^2, \\
Q_0(\tau) &= 1 - (r_1 + r_2)\tau = 1 + (s - s^{-1})\tau, \\
Q_1(\tau) &= \tau
\end{aligned}$$

and we obtain

$$\mathcal{AC}_n(t_1, \ldots, t_k) = q(t_1)^{-1} \ldots q(t_k)^{-1} \sum_{0 \le j_1, \ldots, j_k \le 1} Q_{j_1}(t_1) \ldots Q_{j_k}(t_k) \nabla_n(x_{i_1}^{j_1} \ldots x_{i_k}^{j_k}).$$

□

Using the recurrence relation we give some examples of Alexander-Conway polynomials:

Example 3.1. The recurrence relation for $\nabla_2(x_1^a)$ starts with $\nabla_2(1) = 0$, $\nabla_2(x_1) = 1$ and gives the general term,

$$\nabla_2(x_1^a) = \frac{s^{-a} + (-1)^{a+1} s^a}{s + s^{-1}} = s^{1-a} - s^{3-a} + s^{5-a} - \ldots + (-1)^{a+1} s^{a-1},\ a \ge 0,$$

and $\nabla_2(x_1^a) = (-1)^{a+1}s^{a+1} + (-1)^a s^{a+3} + \ldots - s^{-a-3} + s^{-a-1}$, $a < 0$.

Example 3.2. With specializations $(l, m) \longrightarrow (\iota, \iota(s^{-1} - s))$ in Example 2.1, we have recurrences for $\nabla(k) = \nabla_3(\gamma_k)$, γ_k is the 3-braid $x_1x_2x_1\ldots$ (k-factors):

1) $\nabla(2k+1) = (s^{-1} - s)\nabla(2k) + \nabla(2k-1)$
2) $\nabla(6k) = (s^{-1} - s)[\nabla(6k-1) + \nabla(6k-3) + \nabla(6k-5)] + \nabla(6k-6)$
3) $\nabla(6k+2) = (s^{-1} - s)[\nabla(6k+1) + \nabla(6k-1)] + \nabla(6k-2)$
4) $\nabla(6k+4) = (s^{-1} - s)\nabla(6k+3) + \nabla(6k+2)$.

Remark 3.3. From these recurrences we can compute Jordan normal form of the recurrence matrix $M(k)$:

$$M(k) \sim \begin{pmatrix} J_3 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & s^6 & 0 \\ 0 & 0 & 0 & s^{-6} \end{pmatrix}, \text{ where } J_3 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}.$$

Example 3.4. The first six Alexander-Conway polynomials are

$\nabla(0) = 0$
$\nabla(1) = 0$
$\nabla(2) = 1$
$\nabla(3) = s^{-1} - s$
$\nabla(4) = s^{-2} - 1 + s^2$
$\nabla(5) = s^{-3} - s^{-1} + s - s^3$

Using recurrences for $\nabla(k)$ in Example 3.2, we have explicit formulae:

Proposition 3.5. $\nabla(6k) = (s^{-4} - s^{-2})\sum_{i=0}^{k-1} s^{-6i} + (s^4 - s^2)\sum_{i=0}^{k-1} s^{6i}$

$$\nabla(6k+1) = (s^{-5} - s^{-3})\sum_{i=1}^{k-1} s^{-6i} - (s^5 - s^3)\sum_{i=0}^{k-1} s^{6i}$$

$$\nabla(6k+2) = (s^{-6} - s^{-4})\sum_{i=0}^{k-1} s^{-6i} + 1 + (s^6 - s^4)\sum_{i=0}^{k-1} s^{6i}$$

$$\nabla(6k+3) = (s^{-7} - s^{-5})\sum_{i=0}^{k-1} s^{-6i} + s^{-1} - s - (s^7 - s^5)\sum_{i=0}^{k-1} s^{6i}$$

$$\nabla(6k+4) = (s^{-2} - 1)\sum_{i=0}^{k} s^{-6i} + 1 + (s^2 - 1)\sum_{i=0}^{k} s^{6i}$$

$$\nabla(6k+5) = (s^{-3} - s^{-1})\sum_{i=0}^{k} s^{-6i} - (s^3 - s^1)\sum_{i=0}^{k} s^{6i}$$

In the first four relations k is positive, in the last two, k is nonnegative.

Definition 3.6. For a non zero Laurent polynomial $P = a_p s^p + a_{p-1}s^{p-1} + \ldots + a_{q+1}s^{q+1} + a_q s^q \in \mathbb{C}[s, s^{-1}]$, where p and q are integers and $a_p \neq 0$, $a_q \neq 0$, $p \geq q$, p

is called the *degree of* P, $\deg(P)$, q is called the *order of* P, $\mathrm{ord}(P)$ and $p-q+1$ is called the *breadth of* P, $\mathrm{br}(P)$. For $p=0$ we take $\deg(0)=-\infty, \mathrm{br}(0)=0$.

Proposition 3.7. *If* $a_i \in \{0,1\}$, *then*

$$\deg\nabla_3(\underbrace{x_1^{a_1}x_2^{a_2}\dots x_1^{a_{2k-1}}x_2^{a_{2k}}}_{2k\,factors}) \le 2k-2.$$

Proof. We prove the result by induction: for $k=1$,

$$\nabla(x_1^0x_2^0)=\nabla(x_1^1x_2^0)=\nabla(x_1^0x_2^1)=0,\ \nabla(x_1x_2)=1.$$

Suppose that the result is true for $k-1$, i.e $\deg\nabla_3(x_1^{a_1}x_2^{a_2}\dots x_1^{a_{2k-3}}x_2^{a_{2k-2}}) \le 2k-4$.

For $\beta = x_1^{a_1}x_2^{a_2}\dots x_2^{a_{2k-2}}x_1^{a_{2k-1}}x_2^{a_{2k}}$, we have two cases:

Case I: If $a_1=a_2=\dots=a_{2k}=1$, we have equality (see Proposition 3.5).

Case II: If there is at least one zero, say $a_{2k}=0$, then β is conjugate to $x_1^{a_1+a_{2k-1}}x_2^{a_2}\dots x_1^{a_{2k-3}}x_2^{a_{2k-2}}$. If $a_1'=a_1+a_{2k-1}\le 1$, then it is clear using induction step, otherwise use expansion formula

$$\begin{aligned}\nabla_3(x_1^{a_1}x_2^{a_2}\dots x_1^{a_{2k-1}}x_2^{a_{2k}}) &= \nabla_3(x_1^2x_2^{a_2}\dots x_2^{a_{2k-2}})\\ &= \left(\frac{s}{s^2+1}\right)^{2k-2}\sum_{0\le j_*\le 1} C^{2+j_1}\dots C^{a_{2k-2}+j_{2k-2}}\nabla_3(x_1^{j_1}\dots x_2^{j_{2k-2}})\end{aligned}$$

$\deg(\frac{s}{s^2+1})^{2k-2} = -(2k-2)$. As $a_i+j_i\in\{0,1,2\}$ for $i=2,\dots,2k-2$ and $C^0=C^2=s+s^{-1}$, $C^1=0$, $C^3=-s^2+s^{-2}$
$\max\{\deg(C^{2+j_1}\dots C^{a_{2k-2}+j_{2k-2}})$, where $0\le j_1,j_2,\dots,j_{2k-2}\le 1\}$
$=2+\max\{\deg(C^{a_2+j_2}\dots C^{a_{2k-2}+j_{2k-2}})$, where $0\le j_2,\dots,j_{2k-2}\le 1\}$
$=2+2k-3=2k-1$
Also from induction hypothesis $\deg\nabla_3(x_1^{j_1}\dots x_2^{j_{2k-2}})\le 2k-4$. So

$$\deg\nabla_3(x_1^{a_1}x_2^{a_2}\dots x_1^{a_{2k-1}}x_2^{a_{2k}}) \le -(2k-2)+2k-1+2k-4 \le 2k-2.$$

□

For large exponents, we have

Theorem 3.8. *If all* $a_i\ge 2$, *then*

$$\deg\nabla_3(x_1^{a_1}x_2^{a_2}\dots x_1^{a_{2k-1}}x_2^{a_{2k}}) = a_1+a_2+\dots+a_{2k}-2.$$

Proof. To evaluate $\nabla_3(x_1^{a_1}x_2^{a_2}\dots x_1^{a_{2k-1}}x_2^{a_{2k}})$, use the expansion formula and we have

$$\nabla_3(x_1^{a_1}x_2^{a_2}\,..\,x_1^{a_{2k-1}}x_2^{a_{2k}}) \;=\; \left(\frac{s}{s^2+1}\right)^{2k}\sum_{0\le j_*\le 1} C^{a_1+j_1}\,..\,C^{a_{2k}+j_{2k}}\nabla_3(x_1^{j_1}\,..\,x_2^{j_{2k}})$$

$\deg(\frac{s}{s^2+1})^{2k} = -2k$, Proposition 3.7 and Proposition 3.5 imply that

$$\deg\nabla_3(\underbrace{x_1x_2x_1x_2\dots x_1x_2}_{2k\,\text{factors}}) = 2k-2 \quad (j_1 = j_2 = \dots j_{2k} = 1)$$

As $\deg C^{a_i+1} = a_i > a_i - 1 = \deg C^{a_i+0}$, so
$\max\{\deg(C^{a_1+j_1}C^{a_2+j_2}\dots C^{a_{2k}+j_{2k}})$, where $0 \le j_1, j_2, \dots, j_{2k} \le 1\}$
$= \deg(C^{a_1+1}C^{a_2+1}\dots C^{a_{2k}+1}) = a_1 + a_2 + \dots + a_{2k}$.
Finally

$$\begin{aligned}\deg\nabla_3(x_1^{a_1}x_2^{a_2}\dots x_1^{a_{2k-1}}x_2^{a_{2k}}) &= -2k + a_1 + a_2 + \dots + a_{2k} + 2k - 2\\ &= a_1 + a_2 + \dots + a_{2k} - 2.\end{aligned}$$

□

Corollary 3.9. *If all $a_i \ge 2$, then*

$$br(\nabla_3(x_1^{a_1}x_2^{a_2}\dots x_1^{a_{2k-1}}x_2^{a_{2k}})) = 2(a_1 + a_2 + \dots + a_{2k}) - 3.$$

4. **The degenerate case**

The specialization $l = s, m = -2$, corresponding to the roots $r_1 = r_2 = s$, gives *the degenerate polynomial D* with skein relation

$$sD(L_+) + s^{-1}D(L_-) - 2D(L_0) = 0$$

and recurrence relation

$$D_n(a_1, \dots, a_j + 2, \dots, a_k) = 2sD_n(a_1, \dots, a_j + 1, \dots, a_k) - s^2D(a_1, \dots, a_j, \dots, a_k).$$

Proof of Theorem 1.8: As in [3] for Jones polynomial and as in section 3 for Alexander-Conway polynomial we find the basic polynomials

$$\begin{aligned}R_0^{[a]}(s) &= -r_1^{a-1}r_2 - r_1^{a-2}r_2^2 - \dots - r_1r_2^{a-1} = (1-a)s^a\\ R_1^{[a]}(s) &= r_1^{a-1} + r_1^{a-2}r_2 - \dots + r_2^{a-1} = as^{a-1}\end{aligned}$$

Using the notations and computations of [9], Remark 2.2, we obtain the result.

□

For a fixed sequence $(x_{i_1}x_{i_2}\dots x_{i_k})$ in $\mathcal{B}_n$ the *generating function* for the degenerate polynomial is defined by the formal series in $t_1,\dots,t_k$:

$$\mathcal{D}_n(t_1,\dots,t_k)=\sum_{(a_1,a_2,\dots,a_k)\in\mathbb{Z}^k} D_n(x_{i_1}^{a_1}\dots x_{i_k}^{a_k})(s)t_1^{a_1}\dots t_k^{a_k}.$$

Proof of Theorem 1.10: Using [9], Theorem 3.8, the corresponding polynomials are $e(\tau)=(1-s\tau)^2$ and $E_0(\tau)=1-2s\tau$, $E_1(\tau)=\tau$.

□

Proposition 4.1. ***(Properties of** D **polynomial)*** *Let L, L_1, L_2 be oriented links, $\overline{L}$ be the mirror image of L, $\overleftarrow{L}$ the link with reversed orientation of every component of $\overrightarrow{L}$, "$\coprod$" be the distant union of two links, then*

1) $D(L_1\coprod L_2)=\frac{1}{2s}(s^2+1)D(L_1)D(L_2)$
2) $D(\overline{L})(s)=D(L)(s^{-1})$
3) $D(L_1+L_2)=D(L_1)D(L_2)$
4) $D(\overleftarrow{L})=D(\overrightarrow{L})$.

Proof. The proofs are as in [8] for Jones polynomial. □

Remark 4.2. a) For any link L, $D(L)(1)=1$.
b) The degenerate polynomials take values in $\mathbb{Z}[\frac{1}{2}][s,s^{-1}]$.

Example 4.3. Starting with $D(\bigcirc)=D_2(x_1)=1$ we obtain $D(\bigcirc\bigcirc)=D_2(1)=\frac{1}{2}(s+s^{-1})$ and $D_2(x_1^n)=\frac{1}{2}[(1-n)s^{n+1}+(1+n)s^{n-1}]$; in particular $D_2(x_1^2)=\frac{1}{2}(-s^3+3s)$ and $D_2(x_1^3)=-s^4+2s^2$.

Example 4.4. With specializations $(l,m)\longrightarrow(s,-2)$ in Example 2.1, we have the following recurrences for $D(k)=D_3(\gamma_k)$ (γ_k is the 3-braid $x_1x_2x_1\dots$ with $k-$ factors):

1) $D(2k+1)=2sD(2k)-s^2D(2k-1)$
2) $D(6k+4)=2sD(6k+3)-s^2D(6k+2)$
3) $D(6k+2)=2sD(6k+1)-2s^3D(6k-1)+s^4D(6k-2)$
4) $D(6k)\quad=2sD(6k-1)-2s^3D(6k-3)+2s^5D(6k-5)-s^6D(6k-6)$.

Remark 4.5. These recurrences relations for D polynomials give a recurrence matrix with Jordan normal form:

$$M(k)\sim\begin{pmatrix} J_3 & 0 & 0 & 0\\ 0 & s^6 & 0 & 0\\ 0 & 0 & \gamma s^6 & 0\\ 0 & 0 & 0 & \overline{\gamma}s^6\end{pmatrix},\text{ with } J_3=\begin{pmatrix}0&1&0\\0&0&1\\0&0&0\end{pmatrix},\ \gamma=4+\iota\sqrt{3}.$$

Example 4.6. First few $D(k) = D_3(\gamma_k)$ polynomials are:
$D(0)=\frac{1}{4}(s^2+2+s^{-2})$, $D(1) = \frac{1}{2}(s+s^{-1})$, $D(2) = 1$, $D(3) = \frac{1}{2}(-s^3+3s)$, $D(4) = -s^4+2s^2$, $D(5) = \frac{1}{2}(-3s^5+5s^3)$, $D(6) = \frac{1}{4}(-s^8-6s^6+11)$ and for $n \geq 7, D(n) = \frac{1}{4}(-a_n s^{n+2} + b_n s^{n+1} + c_n s^n)$, where the sequence $(a_n)_{n\geq 6}$ is given by $a_{12k} = 12k-1$, $a_{12k+i} = 12k$ for $i = 1,\dots,5$, $a_{12k+6} = 12k+1$, $a_{12k+6+i} = 12k+2i$ for $i = 1,\dots,5$.

Example 4.7. If $\gamma = \alpha x_n^k \beta$ is an $n+1$-braid with $\alpha, \beta \in \mathcal{B}_n$, then

$D_{n+1}(\gamma) = D_n(\alpha\beta) D_2(x_1^k)$. In particular $D_3(x_1^a x_2^b) = D_2(x_1^a) D_2(x_1^b)$.

Proposition 4.8. *If* $a_i \geq 0$ *and the* 3*-braid* $x_1^{a_1} \dots x_2^{a_{2k}}$ *has degree* $A = \sum_{i=1}^{2k} a_i$ *then*

$$deg D_3(x_1^{a_1} x_2^{a_2} \dots x_1^{a_{2k-1}} x_2^{a_{2k}}) \leq A + 2.$$

Proof. By induction on k and by previous example, this inequality is true for exponents $0 \leq a_i \leq 1$. For arbitrary non negative exponents the general expansion formula gives

$$D_3(a_1, \dots, a_{2k}) = \sum_{J_* \in \{0,1\}^{2k}} R_{J_*}^{A_*} D_3(j_1, j_2, \dots, j_{2k}),$$

where $R_{J_*}^{A_*} = R_{j_1}^{[a_1]} \dots R_{j_k}^{[a_{2k}]}$ has the degree at most $\sum_{i=1}^{2k}(a_i - j_i) = A - J$ (zero occurs only if $(a_i, j_i) = (0,1) \,\text{or}\, (1,0)$) and $D_3(j_1, \dots, j_{2k})$ has degree $\leq J+2$. ☐

Example 4.9. $D_3(x_1^{a_1} x_2^{a_2}) = \frac{1}{4}(1-a_1)(1-a_2)s^{A+2} + \dots$

Example 4.10. $D_3(x_1^{a_1} x_2^{a_2} x_1^{a_3} x_2^{a_4}) = \frac{1}{4}[1 - A + (a_1+a_3)(a_2+a_4) - a_1a_2a_3a_4]s^{A+2} + \dots$

Remark 4.11. In these cases ($k = 1$ and $k = 2$), for exponents $a_i \geq 2$, we have equality in Proposition 4.8. Perhaps the same is true for arbitrary k.

5. HOMFLY polynomial for simple braids

We describe a subset of divisors of Garside braid Δ_n which is related with HOMFLY recurrence and also has nice properties with respect to conjugation in $\mathcal{B}_n$. We denote by $\mathcal{MB}_n$ the monoid of positive braids and by $\text{Div}(\Delta_n)$ the set of subwords of Δ_n (which coincides with the set of left divisors, the set of right divisors of Δ_n, and also with the set of square free elements of $\mathcal{MB}_n$, see [7]). It is also well known that conjugation of positive braids α, α' in $\mathcal{B}_n$ is equivalent with conjugation in $\mathcal{MB}_n$: there is a positive braid β such that $\alpha\beta = \beta\alpha'$. Initial terms in the recurrence relation for HOMFLY are polynomials $P(\widehat{\beta}) = P_n(\beta)$ for positive braids which have, up to a conjugation, all exponents equal to 1. One can define this subset, denoted by $\mathcal{SB}_n$, of positive braids in many ways:

Definition 5.1. $\beta \in \mathcal{MB}_n$ is *simple* if one presentation of β contains no repeated generators x_i (in this case, any presentation of β as a positive braid has the same property).

Definition 5.2. $\beta \in \mathcal{MB}_n$ is *simple* if all its conjugates in $\mathcal{MB}_n$ are square free.

The equivalence of these definitions is proved in [2]. In the same paper one can find canonical forms for simple braids as a product of disjoint cycles and the canonical form for conjugacy class of a simple braid (see Theorem 1.11 a)). Here is a picture of a simple braid: if $A = (4, 3, 2, 2)$ and $\beta_A = (x_1x_2x_3)(x_5x_6)(x_8)(x_{10})$ is in $\mathcal{B}_{13}$, then



It is obvious that the closed braid $\widehat{\beta_A}$ is the trivial link with $n-s_r+r = n-\deg(\beta)$ components.

Theorem 5.3. *The value of HOMFLY polynomial $P_n(\beta)$ of a simple braid β is:*

$$P_n(\beta) = \Big(-\frac{l+l^{-1}}{m}\Big)^{n-deg(\beta)-1}.$$

Proof. The closure of the simple braid β is the trivial link with $n - \deg(\beta)$ components and $P_n(\text{unlink with } k \text{ components}) = \Big(-\frac{l+l^{-1}}{m}\Big)^{k-1}$. Alternatively, one can use Markov moves $d = \deg(\beta)$ times and obtain $P_n(\beta) = P_{n-d}(1)$. □

Proof of Theorem 1.11 b): The degree of the simple braid

$$\beta_A = (x_1x_2 \dots x_{s_1-1})(x_{s_1+1} \dots x_{s_2-1}) \dots (x_{s_{r-1}+1} \dots x_{s_r-1})$$

is $s_r - r$.

□

With specializations $(l, m) \longrightarrow (\iota, \iota(s^{-1} - s)), (l, m) \longrightarrow (\iota s^2, \iota(s - s^{-1}))$ and $(l, m) \longrightarrow (s, -2)$ respectively, we obtain the initial values for recurrence:

Corollary 5.4. $\nabla_n(\beta_A) = \begin{cases} 0, \text{ if } deg(\beta) \neq n-1 \\ 1, \text{ if } deg(\beta) = n-1. \end{cases}$

Corollary 5.5. $V_n(\beta_A) = (-\frac{s^2+1}{s})^{n-deg(\beta)-1}$.

Corollary 5.6. $D_n(\beta_A) = \Big(\frac{s^2+1}{2s}\Big)^{n-deg(\beta)-1}$.

6. Relation between link polynomials

Using previous computations for some families of braids (Examples 3.1, 4.3, 4.6 and Corollaries 5.5 and 5.6) we can give "theoretical " proofs of the independence of the link polynomials ∇, V and D. More important for knot theorist is an example where D can differentiate links and one of the links of [5] is good for this.

Definition 6.1. A set of k families of Laurent polynomials $\{f_1^\alpha, f_2^\alpha, \dots, f_k^\alpha\}_{\alpha\in\mathcal{A}},\ f_i^\alpha \in \mathbb{C}[s,s^{-1}]$ is called *linearly independent* over $\mathbb{C}[s,s^{-1}]$ if from an equation

$$\sum_{i=1}^{k} g_i(s,s^{-1}) f_i^\alpha = 0, \ \ (g_i(s,s^{-1}) \in \mathbb{C}[s,s^{-1}])$$

which is satisfied for any value of parameter α in $\mathcal{A}$ we obtain $g_i = 0$, $i = 1,\dots,k$.

Definition 6.2. A set of k families of Laurent polynomials $\{f_1^\alpha, \dots, f_k^\alpha\}_{\alpha\in\mathcal{A}}$, $f_i^\alpha \in \mathbb{C}[s,s^{-1}]$ are called *algebraically independent* over $\mathbb{C}[s,s^{-1}]$ if from a polynomial equation $\Phi(f_1^\alpha,\dots,f_k^\alpha) = 0$, with $\Phi \in \mathbb{C}[s,s^{-1}][T_1,\dots,T_k]$, which is satisfied for any $\alpha \in \mathcal{A}$, we obtain $\Phi = 0$.

As a first example we take $\mathcal{A} = \mathcal{B}_2$, $f_1(\beta) = \nabla_2(\beta)$, $f_2(\beta) = V_2(\beta)$, $f_3(\beta) = D_2(\beta)$ and we obtain

Proposition 6.3. *$\{\nabla_2(\beta), V_2(\beta), D_2(\beta)\}_{\beta\in\mathcal{B}_2}$ is a linearly independent family over $\mathbb{C}[s,s^{-1}]$.*

Proof. Suppose we have relations $A\nabla_2(x_1^n)+BV_2(x_1^n)+CD_2(x_1^n) = 0$ where $A, B, C \in \mathbb{C}[s,s^{-1}]$ and $\deg V_2(x_1^n) = 3n-1$, $\deg\nabla_2(x_1^n) = n-1$, $\deg D_2(x_1^n) = n+1$ $(n \geq 2)$. If B in not zero, take $b = \deg B$ and choose n big enough to have $b + (3n-1) > \max\{\deg A, \deg C\} + n + 1$. Then the equation

$$A(s,s^{-1})V_2(x_1^n) + B(s,s^{-1})\nabla_2(x_1^n) + C(s,s^{-1})D_2(x_1^n) = 0$$

gives a contradiction. Therefore $B = 0$ and for A, C non zero we find a contradiction comparing the order of $\nabla_2(x_1^n)$ (which is $1-n$) and the order of $D_2(x_1^n)$, which is positive. □

In the second example we choose $\mathcal{A} = \mathcal{SB} = \coprod\limits_n \mathcal{SB}_n$ the set of positive braids, $f_1(\beta) = V(\widehat{\beta})$ and $f_2(\beta) = D(\widehat{\beta})$.

Proposition 6.4. *$\{V(\widehat{\beta}), D(\widehat{\beta})\}_{\beta\in\mathcal{SB}}$ is an algebraic independent family over $\mathbb{C}[s,s^{-1}]$.*

Proof. Take $P(Y,Z)$ a polynomial with coefficients in $\mathbb{C}[s,s^{-1}]$ which is zero for $Y = V(\widehat{\beta}), Z = D(\widehat{\beta})$ and any simple braid β. We will show that P is identically zero. Looking at the family $\beta_k^n = x_1x_2\dots x_{k-1}$, $2 \leq k \leq n-1$ and $k \equiv n \mod (2)$, we find $V(\widehat{\beta_k^n}) = (\frac{s^2+1}{s})^{n-k}$, and $D(\widehat{\beta_k^n}) = (\frac{s^2+1}{2s})^{n-k}$ and also $P(V(\widehat{\beta_k^n}), D(\widehat{\beta_k^n})) = 0$.

Now we will prove that top degree monomials of P are zero :

$$P(Y,Z)=\sum_{i=0}^{d}A_{d-i,i}(s,s^{-1})Y^{d-i}Z^{i}+\sum_{0\leq i+j\leq d-1}A_{i,j}(s,s^{-1})Y^{i}Z^{j}$$

For $Y=V(\widehat{\beta_k^n})$ and $Z=D(\widehat{\beta_k^n})$ we have

$$\Big[\sum_{i=0}^{d}\frac{1}{2^{(n-k)i}}A_{d-i,i}(s,s^{-1})\Big]\Big(\frac{s^2+1}{s}\Big)^{d(n-k)}+\ldots=0$$

and for $n-k$ big enough we have

$$\max(\deg A_{d-i,i})+d(n-k)>\max(\deg A_{i,j})+(d-1)(n-k),$$

therefore Laurent polynomial $\sum\limits_{i=0}^{d}\frac{1}{2^{(n-k)i}}A_{d-i,i}(s,s^{-1})$ should be zero. Induction on i ends the proof : for $n-k\to\infty$ we obtain $A_{d,0}(s,s^{-1})=0$, and if $A_{d,0}=A_{d-1,1}=\ldots=A_{d-i+1,i-1}=0$, then $2^{i(n-k)}[\frac{1}{2^{i(n-k)}}A_{d-i,i}(s,s^{-1})+\frac{1}{2^{(i+1)(n-k)}}A_{d-i-1,i+1}(s,s^{-1})+\ldots]$ is zero and $n-k\to\infty$ gives $A_{d-i,i}(s,s^{-1})=0$. $\square$

As a final example we take $\mathcal{A}=\{\text{isotopy classes of links}\}$, $f_1(L)=\nabla(L)$, $f_2(L)=V(L)$ and $f_3(L)=D(L)$ and we show that $\{\nabla(L),V(L),D(L)\}_{\mathcal{A}}$ is algebraically independent over $\mathbb{C}[s,s^{-1}]$.

Proof of Theorem 1.12 Take p a prime number, n an even natural number and the link $L_{p,n}=\widehat{\beta}_{p,n}$, where $\beta_{p,n}$ is the $n+1$-braid $x_1^{2p+1}x_2^{2p+1}\ldots x_n^{2p+1}$. The corresponding polynomials of $L_{p,n}$ have leading terms $\nabla(L_{p,n})=(s^{2p}+\ldots)^n$, $V(L_{p,n})=(s^{6p+2}+\ldots)^n$, and $D(L_{p,n})=(ps^{2p+2}+\ldots)^n$. Suppose that

$$P(X,Y,Z)=\sum_{i,j,k}A_{i,j,k}(s,s^{-1})X^iY^jZ^k$$

is a nonzero polynomial with coefficients in $\mathbb{C}[s,s^{-1}]$ which is zero for $(X,Y,Z)=(\nabla(L_{p,n}),V(L_{p,n}),D(L_{p,n}))$ (only nonzero coefficient $A_{i,j,k}$ are in this sum, $\alpha_{i,j,k}s^{a_{i,j,k}}$is the leading coefficient of $A_{i,j,k}(s,s^{-1})$). We will find a contradiction if P is nonzero. The leading term of the polynomial in s $P(L_{p,n})=P(\nabla(L_{p,n}),V(L_{p,n}),D(L_{p,n}))$ is given by

$$\sum\alpha_{i,j,k}p^{nk}s^{a_{i,j,k}+2pni+(6p+2)nj+(2p+2)nk}$$

where $a_{i,j,k}+2pni+(6p+2)nj+(2p+2)nk$ is maximal (i,j,k are variable). Its coefficient $\sum\limits_{i,j,k}\alpha_{i,j,k}p^{nk}$ should be zero, and the polynomial equation (complex coefficients) $\sum\limits_{i,j,k}\alpha_{i,j,k}t^k=0$ cannot have infinitely many solutions (p^n), therefore for a fixed k, $\sum\limits_{i,j}\alpha_{i,j,k}=0$. From maximality condition i and j are such that the sum $S_{i,j} = a_{i,j,k} + 2pni + (bp + 2)nj$ is constant in i and j (for any p and n); we show that this implies the uniqueness of the pair (i, j), hence we find a contradiction: $\alpha_{i,j,k} = 0$. First choose n greater than any difference $|a_{i,j,k} - a_{h,l,k}|$, next choose p greater than any difference $|a_{i,j,k} - a_{h,l,k}|$ and $|a_{i,j,k} + 2nj - a_{h,l,k} - 2nl|$ (condition on n shows that the last difference is not 0 for $j \neq l$). If there are two pairs $(i, j), (h, j)$ with $S_{i,j} = S_{h,j}$, we find $a_{i,j,k} + 2pni = a_{h,j,k} + 2pnh$, hence $p \,|\, (a_{i,j,k} - a_{h,j,k})$ and this is possible only for $a_{i,j,k} = a_{h,j,k}$, and the previous equation gives $i = h$. If there are two pairs $(i, j), (h, l)$ $(j \neq l)$ with $S_{i,j} = S_{h,l}$, we find $a_{i,j,k} + 2pni + (6p+2)nj = a_{h,l,k} + 2pnh + (6p+2)nl$, hence $p \,|\, (a_{i,j,k} + 2nj - a_{h,l,k} - 2nl)$ and this is not possible because $0 < |a_{i,j,k} + 2nj - a_{h,l,k} - 2nl| < p$.

$\square$

Example 6.5. (Eliahou, Kauffman, Thislethwaite [5]) The link $LL_2(2)$ has the same Jones and Alexander-Conway polynomials, and different D polynomials. Using the relative expansion formula $\mathcal{L}(-5, 6)$ we have to compute $D_{\mathcal{L}}(0, 0)$, $D_{\mathcal{L}}(0, 1)$, $D_{\mathcal{L}}(1, 0)$ and $D_{\mathcal{L}}(1, 1)$:

$$D_{\mathcal{L}}(-5, 6) = -30sD_{\mathcal{L}}(0, 0) + 36D_{\mathcal{L}}(0, 1) + 25D_{\mathcal{L}}(1, 0) - 30s^{-1}D_{\mathcal{L}}(1, 1).$$

$\mathcal{L}(1, 0)$ is the unlink $\bigcirc\bigcirc\bigcirc$, $\mathcal{L}(0, 0)$ is $\text{Hopf}^- \coprod \bigcirc$ and $\mathcal{L}(1, 1)$ is $\text{Hopf}^+ \coprod \bigcirc$ where $\text{Hopf}^- = \widehat{x_1^2}$ and $\text{Hopf}^+ = \widehat{x_1^{-2}}$. $\mathcal{L}(0, 1)$ is a four component link: after two skein decompositions we find $\text{Hopf}^+ \coprod \bigcirc$, the unlink $\bigcirc \bigcirc \bigcirc$ and closure of the 4-braid $x_1^{-1}x_2^2x_1^{-1}x_3^{-3}x_2^2x_3$. Computing only the leading terms, the main contribution comes from the first link and we have $D_{\mathcal{L}}(-5, 6) = \frac{15}{2}s^5 + \ldots$ and $D(\bigcirc\bigcirc) = \frac{s^2+1}{2s}$.

Acknowledgements: The first author has presented a preliminary version of this paper at the Knot Theory Conference held at ICTP in May, 2009. She is grateful to ICTP for inviting her and giving the opportunity to present these results and also to Professor Hugh Morton and Professor Jozef Przytycki for their valuable suggestions they gave during that conference. Both authors are thankful to Pakistan Higher Education Commission for supporting this research.

REFERENCES

[1] E. Artin, *Theory of braids*, Ann. of Math. (2) **48** (1947), 101-126.

[2] R. Ashraf, B. Berceanu : *Simple braids*, in preparation.

[3] B. Berceanu, A.R. Nizami : *Recurrence relation for Jones polynomials*, arXiv:1002.3735v1 (2010).

[4] J. Birman, *Braids, links, and mapping class groups*, Ann. of Math. Studies No. **82**, Princeton University Press, 1974.

[5] S. Eliahou, L.H. Kauffman, M.B. Thislethwaite, *Infinite families of links with trivial Jones polynomial*, Topology 42 (2003), 155-169.

[6] P. Freyd, D. Yetter, J. Hoste, W.B.R. Lickorish, K. Millet and A. Ocneanu : *A new polynomial invariant of knots and links*, Bull.Amer.Math.Soc. 12(1985) 239-246.

[7] F.A. Garside, *The braid groups and other groups*, Quat. J. of Math. Oxford 2^e ser. **20** (1969), 235-254.
[8] W.B.R. Lickorish, *An Introduction to Knot Theory*, Graduate Texts in Mathematics **175**, Springer-Verlag New York, 1997.
[9] A.R. Nizami, *Fibonacci modules and multiple Fibonacci sequences*, arXiv:0810.4028vI (2008), to appear in ARS Combinatoria.

[1] ABDUS SALAM SCHOOL OF MATHEMATICAL SCIENCES, GC UNIVERSITY, LAHORE-PAKISTAN.
E-mail address: `rashraf@sms.edu.pk`

[2] INSTITUTE OF MATHEMATICS SIMION STOILOW, BUCHAREST-ROMANIA (PERMANENT ADDRESS).
E-mail address: `Barbu.Berceanu@imar.ro`